



Complete Summary

TITLE

Thromboprophylaxis: percentage of moderate to high-risk patients (as per guidelines) over 40 years who undergo pelvic floor surgery who receive thromboprophylaxis, during the 6 month time period.

SOURCE(S)

Australian Council on Healthcare Standards (ACHS). ACHS clinical indicator users' manual 2009. ULTIMO NSW: Australian Council on Healthcare Standards (ACHS); 2009 Jan. 853 p.

Measure Domain

PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the [Measure Validity](#) page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This measure is used to assess the percentage of moderate to high-risk patients (as per guidelines*) over 40 years who undergo pelvic floor surgery who receive thromboprophylaxis, during the 6 month time period.

*Refer to the "The Australian and New Zealand Working Party on the Management and Prevention of Venous Thromboembolism. Prevention of Venous Thromboembolism, Best Practice Guidelines for Australia and New Zealand," 3rd Edition, July 2005.

RATIONALE

This indicator has been included as an index of utilisation of evidence-based practice guidelines for thromboprophylaxis in moderate to high-risk women having gynaecological surgery.

PRIMARY CLINICAL COMPONENT

Thromboprophylaxis; pelvic floor surgery; moderate and high-risk women

DENOMINATOR DESCRIPTION

Total number of moderate to high-risk patients (as per guidelines) over 40 years who undergo pelvic floor surgery, during the 6 month time period (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary)

NUMERATOR DESCRIPTION

Total number of moderate to high-risk patients (as per guidelines) over 40 years who undergo pelvic floor surgery who receive thromboprophylaxis, during the 6 month time period

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

- A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence
- A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Use of this measure to improve performance
Variation in quality for the performance measured

EVIDENCE SUPPORTING NEED FOR THE MEASURE

Australian Council on Healthcare Standards (ACHS). Australasian clinical indicator report 2001-2007. Determining the potential to improve quality of care: 9th edition. ULTIMO NSW: Australian Council on Healthcare Standards (ACHS); 2008. 611 p.

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Hospitals

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Single Health Care Delivery Organizations

TARGET POPULATION AGE

Age greater than 40 years

TARGET POPULATION GENDER

Female (only)

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Unspecified

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Unspecified

UTILIZATION

Unspecified

COSTS

Unspecified

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

Moderate to high-risk patients (as per guidelines) over 40 years who undergo pelvic floor surgery, during the 6 month time period

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

Total number of moderate to high-risk patients (as per guidelines*) over 40 years who undergo pelvic floor surgery**, during the 6 month time period

*The definition of **moderate to high-risk** will be taken from "The Australian and New Zealand Working Party on the Management and Prevention of Venous Thromboembolism. Prevention of Venous Thromboembolism, Best Practice Guidelines for Australia and New Zealand," 3rd Edition, July 2005.

- **Moderate risk:**
 - Major surgery, age 40 to 60 years without additional risk factors
 - Minor surgery, age greater than 60 years
 - Minor surgery, age 40 to 60 years with history of venous thromboembolism (VTE) or on oestrogen therapy or with other risk factors
- **High risk:**
 - Orthopedic surgery of pelvis, hip or lower limb
 - Multiple trauma
 - Major surgery, age greater than 60 years
 - Major surgery, age 40 to 60 years with cancer or history of VTE
 - Other risk factors

**Refer to the original measure documentation for specific International Classification of Diseases, Tenth Revision, Australian Modification (ICD-10-AM) codes.

Exclusions

Unspecified

RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

DENOMINATOR (INDEX) EVENT

Clinical Condition
Institutionalization
Therapeutic Intervention

DENOMINATOR TIME WINDOW

Time window brackets index event

NUMERATOR INCLUSIONS/EXCLUSIONS**Inclusions**

Total number of moderate to high-risk patients (as per guidelines*) over 40 years who undergo pelvic floor surgery who receive thromboprophylaxis, during the 6 month time period

*See the "Denominator Inclusions/Exclusions" field.

Exclusions

Unspecified

MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Administrative data
Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

External comparison at a point in time

External comparison of time trends

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Indicator area 7: thromboprophylaxis CI 7.2.

MEASURE COLLECTION

[Australian Council on Healthcare Standards \(ACHS\) Equip Clinical Indicators](#)

MEASURE SET NAME

[Gynaecology Indicators](#)

DEVELOPER

Australian Council on Healthcare Standards

FUNDING SOURCE(S)

Funding is direct Australian Council on Healthcare Standards (ACHS) funding sourced through our membership. ACHS does not receive external funding from the government or other sources.

COMPOSITION OF THE GROUP THAT DEVELOPED THE MEASURE

Our terms of reference dictate the composition of the working parties that develop our indicators and include the following:

- Two Clinicians -- nominated by the relevant specialty college/association/society, one nominated to be the chair of the working party
- Private Hospital Representative -- nominated by the Australian Private Hospital Association
- Consumer Representative -- nominated by the Consumer Health Forum of Australia
- Coding Representative -- nominated by the National Centre for Clinical classification on Health
- Quality Health New Zealand, nominated by QH NZ (if applicable)
- Epidemiological/Clinical Research Representative, Director of Health Services Research Group, University of Newcastle
- Australian Council on Healthcare Standards (ACHS) Representatives -- Clinical Director, Coordinator, Administrative Assistant
- Other Expert Stakeholders, as required

FINANCIAL DISCLOSURES/OTHER POTENTIAL CONFLICTS OF INTEREST

None

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

1995 Jan

REVISION DATE

2009 Jan

MEASURE STATUS

This is the current release of the measure.

This measure updates a previous version: Australian Council on Healthcare Standards (ACHS). ACHS clinical indicator users' manual 2008. ULTIMO NSW: Australian Council on Healthcare Standards (ACHS); 2007 Dec. 776 p.

SOURCE(S)

Australian Council on Healthcare Standards (ACHS). ACHS clinical indicator users' manual 2009. ULTIMO NSW: Australian Council on Healthcare Standards (ACHS); 2009 Jan. 853 p.

MEASURE AVAILABILITY

The individual measure, "Indicator Area 7: Thromboprophylaxis CI 7.2," is published in "ACHS Clinical Indicator Users' Manual 2009."

For more information contact, the Australian Council on Healthcare Standards (ACHS), 5 Macarthur Street, ULTIMO NSW 2007; Phone: (02) 9281 9955; Fax: (02) 9211 9633; E-mail: pos@achs.org.au; Web site: www.achs.org.au.

COMPANION DOCUMENTS

The following is available:

- Australian Council on Healthcare Standards (ACHS). Australasian clinical indicator report 2001-2007. Determining the potential to improve quality of care: 9th edition. ULTIMO NSW: Australian Council on Healthcare Standards (ACHS); 2008. 611 p. This document is available in Portable Document Format (PDF) from the [Australian Council on Healthcare Standards \(ACHS\) Web site](http://www.achs.org.au).

NQMC STATUS

This NQMC summary was completed by ECRI Institute on November 5, 2007. The information was verified by the measure developer on March 10, 2008. This NQMC summary was updated by ECRI Institute on May 8, 2009.

COPYRIGHT STATEMENT

This NQMC summary is based on the original measure, which is subject to the measure developer's copyright restrictions. This work is copyright. Apart from any use as permitted under the Copyright Act 1968, no part may be reproduced by any process without written permission from The Australian Council on Healthcare Standards (ACHS).

Disclaimer

NQMC DISCLAIMER

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at <http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx>.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

[Copyright/Permission Requests](#)

Date Modified: 7/27/2009

